

Heart Failure

| Key Indicators | Trinity Health (1) | National Average (2) | Top 10% of Hospitals Nationwide (2) | Trinity Health Hospitals at Top 10% |
|---|--------------------|----------------------|-------------------------------------|-------------------------------------|
| Percent of Heart Failure Patients Given Discharge Instructions | 91% | 79% | 100% | 2 out of 24 hospitals |
| Percent of Heart Failure Patients Given an Evaluation of Left Ventricular Systolic (LVS) Function | 100% | 90% | 100% | 17 out of 24 hospitals |
| Percent of Heart Failure Patients Given ACE Inhibitor or ARB for Left Ventricular Systolic Dysfunction (LVSD) | 97% | 90% | 100% | 12 out of 24 hospitals |
| Percent of Heart Failure Patients Given Smoking Cessation Advice/Counseling | 100% | 93% | 100% | 24 out of 24 hospitals |

*The percentage includes only patients whose history and condition indicate the treatment is appropriate. Talk to your doctor if you have questions about your treatment.

- (1) Trinity Health data time period: October 2009 to March 2010.
- (2) National average and Top 10% of hospitals nationwide were obtained from the Hospital Compare website on July 21, 2010. The latest reporting period is from October 2008 through September 2009 for all indicators.